

MedChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: C. Schmidt, B. Karge, R. Misgeld, A. Prokop, M. Brönstrup and I. Ott, *Med. Chem. Commun.*, 2017, DOI: 10.1039/C7MD00269F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Biscarbene gold(I) complexes: Structure-activity-relationships regarding antibacterial effects, cytotoxicity, TrxR inhibition and cellular bioavailability

Claudia Schmidt¹, Bianka Karge², Rainer Misgeld³, Aram Prokop³, Mark Brönstrup² and Ingo Ott¹

¹Institute of Medicinal and Pharmaceutical Chemistry, Technische Universität Braunschweig, Beethovenstr. 55, 38106 Braunschweig, Germany; ²Department of Chemical Biology, Helmholtz Centre for Infection Research and German Centre for Infection Research (DZIF) Inhoffenstr. 7, 38124 Braunschweig, Germany; ³Department of Pediatric Oncology, Children's Hospital Cologne, Amsterdamer Strasse 59, 50735 Cologne, Germany

Keywords:

anticancer, antibacterial, bioorganometallics, carbene, gold, inorganic medicinal chemistry, thioredoxin reductase

Abstract

A series of gold(I) complexes with two *N*-heterocyclic carbene ligands (biscarbene gold complexes) was prepared and evaluated for effects against cancer cells and pathogenic bacteria. Proliferation inhibition was observed in cancer cells and in Gram-positive bacteria, whereas Gram-negative bacteria were less sensitive towards the compounds. The protein binding and cellular uptake were quantified and the combined results indicated a strong correlation between cellular bioavailability and antiproliferative effects. The biscarbene gold complexes inhibited bacterial and mammalian TrxRs with low to moderate potency. However, based on the obtained structure-activity-relationships and the high cellular accumulation levels, TrxR inhibition can be considered as a relevant contributor to the cellular pharmacology of biscarbene gold(I) complexes.

Introduction

Gold complexes have attracted major attention in inorganic medicinal chemistry and are currently considered as metallodrug candidates with several possible therapeutic applications including cancer or infectious diseases.¹⁻⁵ The history of gold in medicine traces back thousands of years, and the element had been of high relevance in alchemy.⁶ In modern science, the probably first report on antibacterial effects of gold salts was published by the famous bacteriologist Robert Koch, and some gold complexes are nowadays used therapeutically in the treatment of rheumatoid arthritis.⁷ Moreover, the strong antiproliferative properties of many gold complexes indicate their potential as anticancer agents, that has currently been evaluated in several ongoing clinical trials with Auranofin.

Regarding gold metallodrug design, the element has been most frequently used in the oxidation state +1 with thiolate and phosphane ligands (as present in the lead compound Auranofin, see Figure 1). However, traditional gold drugs face major challenges regarding their stability, as these ligands are readily replaced under physiological conditions.^{1, 2, 8} More recently, organometallic gold complexes have attracted the attention of inorganic medicinal chemists, as they offer options for a more robust coordination of ligands to the gold central atom. Among those, *N*-heterocyclic carbene ligands are especially promising, because antiproliferative effects, apoptosis induction, inhibition of thioredoxin reductase (TrxR), antibacterial properties and other biological effects have been reported for many derivatives.⁹⁻²⁷ Recently, we described a series of chlorido gold(I) NHC complexes with anticancer and antibacterial properties.¹⁷ The complexes were effective inhibitors of both mammalian and bacterial TrxRs. Of particular importance were the Gram-positive directed antibacterial effects, which might be the consequence of the high dependence of these bacteria on an intact Trx/TrxR system. The dual activity against bacterial and mammalian TrxRs, together with the observed cytotoxic and antibacterial effects, provide the rationale to study gold complexes as both cytotoxic and antibacterial agents. In this report, we extend these studies to biscarbene compounds containing two NHC ligands. The cytotoxicity, cellular uptake, antibacterial effects, inhibition of mammalian and bacterial TrxRs, and the resulting structure-activity-relationships are presented.





Figure 1: Auranofin and examples of previously reported gold(I) NHC complexes¹⁷

Chemistry

Starting from the imidazolium cations **1a-1d**, the chlorido gold(I) NHC complexes **2a-2d** were prepared as recently described.¹⁷ The target compounds **3a-3d** were obtained by reacting **2a-2d** with the respective imidazolium cations. All complexes were characterized by ¹H-, ¹³C-NMR, and MS spectroscopy, and high purity of all target compounds was noted by elemental analysis. Mass spectrometry confirmed presence of the respective [M - I]⁺ and [I]⁻ ions and showed the expected isotope pattern in case of the bromine containing compounds. In comparison to the mono-NHC complexes (**2a-2d**), no significant changes could be observed in the ¹H-NMR spectra of **3a-3d**. In the ¹³C-NMR spectra of complexes **3a-d** the signal for the carbon at position 2, which is coordinated to gold(I), is observed at significantly higher ppm values (approx. + 12-13 ppm) than in the mono-NHC precursors **2a-d**.¹⁷



Scheme 1: Synthesis of NHC-Au-NHC complexes.

Cytotoxicity and bacterial growth inhibition

The cytotoxicity and growth inhibition of **3a - 3d** was determined against three cancer cell lines (HT-29 colon carcinoma cells and MCF-7/MDA-MB-231 breast carcinoma cells), one non-tumorigenic kidney cell line (RC-124), as well as seven bacterial strains of the ESKAPE panel (*A. baumannii*, *E. coli*, *K. pneumoniae*, *P. aeruginosa*, *E. faecium*, MRSA RKI, MRSA DSM) in comparison to Auranofin. Regarding the cytotoxicity against cell lines, all complexes were active in the low micromolar to nanomolar range and there was a clear positive effect of the phenyl residues as well as the bromine substituent. Complexes **3c** and **3d** with the phenylimidazole-based NHC ligands had substantially higher activities than Auranofin and reached IC₅₀ values in the range of 0.05 - 0.18 μM. The bromine containing compounds triggered lower IC₅₀ values than the respective bromine free compounds (compare results of the couples **3a / 3b** and **3c / 3d**). In no case a tumor selective toxicity could be reached as the effects against the human embryonic kidney cell line RC-124 were comparable with those against the tumor cell lines.

The most cytotoxic compound **3d** was subjected to further experiments on apoptosis induction in drug resistant p-glycoprotein overexpressing Nalm cells based on our previous

studies with structurally related compounds (e.g. **2b**).^{14, 17} In good agreement with previous observations, **3d** was able to overcome drug resistance in the Daunorubicine and Vincristine-resistant lines (see Figure 2). Importantly, the apoptosis induction was 3-5 fold higher than that of **2b** in the same assay.¹⁷

Compound	HT-29	MCF-7	MDA-MB-231	RC-124
Auranofin	3.79 ^{+/-0.18}	2.00 ^{+/-0.05}	1.54 ^{+/-0.12}	1.44 ^{+/-0.03}
3a	5.54 ^{+/-0.79}	4.20 ^{+/-0.72}	5.08 ^{+/-0.60}	3.44 ^{+/-0.38}
3b	1.51 ^{+/-0.10}	0.72 ^{+/-0.07}	1.08 ^{+/-0.06}	0.94 ^{+/-0.10}
3c	0.15 ^{+/-0.01}	0.16 ^{+/-0.03}	0.18 ^{+/-0.02}	0.09 ^{+/-0.02}
3d	0.14 ^{+/-0.02}	0.06 ^{+/-0.00}	0.18 ^{+/-0.01}	0.05 ^{+/-0.01}

Table 1: Antiproliferative effects of Auranofin and complexes **3a-d** expressed as IC₅₀ values (μM) with standard errors as superscripts.




Figure 2: DNA fragmentation after 72 h of complex **3d** in Nalm-6 and drug resistant Nalm-6 cells. DNR: Daunorubicine, VCR: Vincristin; Nalm-6: wildtype cells; Nalm-6-DNR: Daunorubicine-resistant Nalm-6 cells; Nalm-6-VCR: Vincristin-resistant Nalm-6 cells. K0: untreated control; The IC₅₀ values of **3d** against the different cell lines are: Nalm-6: 0.9 μM, Nalm-6-DNR: 1.3 μM, Nalm-6-VCR: 1.0 μM.

Evaluation of the inhibition of bacterial growth confirmed the previously reported preference for Gram-positive strains (*E. faecium*, MRSA1 and MRSA2) of many gold complexes, as only **3c** and Auranofin triggered moderate activity against Gram-negative bacteria (MIC values of 19.5 μM or higher against *A. baumannii*, *E. coli*, *K. pneumoniae*, *P. aeruginosa*). Against Gram-positive bacteria, the order of activity for the gold NHC complexes was **3c** > **3b** ~ **3d** >

3a, indicating a positive effect of the phenyl substituent. The most potent compound **3c** was in particular active against MRSA with MIC values of approx. 2 μ M.

Compound	MRSA RKI	MRSA DSM	<i>E. faecium</i>	<i>E. coli</i>
3a	>100 (0)	79.0 (29.2)	>100 (0)	>100 (0)
3b	12.7 (3.0)	12.2 (8.0)	70.0 (18.7)	>100 (0)
3c	2.3 (1.2)	1.7 (0.4)	8.7 (0.5)	47.7 (10.7)
3d	16.6 (7.7)	9.6 (3.4)	25.5 (12.6)	>100 (0)
Auranofin	0.6 (0.7)	0.4 (0.3)	0.3 (0.3)	45.7 (5.1)
Antibiotic	2.4 (1.4)	4.7 (5.8)	9.5 (2.0)	0.1 (0)

Compound	<i>P. aeruginosa</i>	<i>A. baumannii</i>	<i>K. pneumoniae</i>
3a	>100 (0)	>100 (0)	>100 (0)
3b	>100 (0)	>100 (0)	>100 (0)
3c	19.5 (6.1)	48.7 (11.0)	85.3 (26.9)
3d	>100 (0)	>100 (0)	>100 (0)
Auranofin	>100 (44.1)	55.0 (18.0)	81.0 (48.0)
Antibiotic	7.3 (1.5)	0.9 (0.6)	0.2 (0.1)

Table 2: Antibacterial activities of **3a-3d** and Auranofin. Minimal inhibitory concentrations (MIC) are given in μ M (standard deviations in brackets); MRSA = Methicillin-resistant *Staphylococcus aureus*. As positive control antibiotics, Amikacin (*P. aeruginosa*), Linezolid (*S. aureus*) and Ciprofloxacin (all other strains) have been used.

Protein Binding and Cellular Uptake

Previous studies on gold NHC complexes had shown correlations between the cytotoxic effects in cells, cellular bioavailability and protein binding.^{15, 17} Protein binding studies of **3a - 3d** with serum albumin and proteins from fetal calf serum in a precipitation assay showed low to moderate binding efficacy. Complex **3c** afforded the highest protein binding (52 % with serum albumin and 56 % with calf serum after 1h), complexes **3a** and **3b** were bound in the rather narrow range of 36 - 47 % after 1h, while **3d** showed the lowest protein binding (8 % with serum albumin and 5 % with calf serum after 1h). The levels remained rather stable over time with only small variations (e.g. some increase in the case of **3d**, see supporting information).

The cellular uptake was quantified in MCF-7 cells, which had been most sensitive towards the complexes (see Table 1). In order to evaluate the influence of protein binding on the bioavailability, the experiments were performed using serum containing and serum free cell

culture media (see Figure 4). However, there was no significant influence of the fetal calf serum proteins on the cellular uptake, as the cellular levels of **3a** - **3d** were comparable with both experimental setups. All compounds showed a fast cellular uptake and reached high values within the first hour of exposure. The highest cellular concentration was noted for complex **3c** (maximum concentration: 9.9 (+FCS) / 9.6 (-FCS) nmol gold/mg cell protein) after 24 h. The most striking result was the substantial difference between the imidazole (**3a,b**) and phenylimidazole (**3c,d**) based gold NHC complexes as the latter ones reached much higher uptake values (2.5 – 7.9-fold higher during 24h). The bromine substituent had a negative impact on the cellular bioavailability as the levels of **3b** and **3d** were within 24 h in most cases up to 4.8-fold lower than those obtained with the respective not bromine-substituted counterparts **3a** and **3c**. Moreover, the levels of **3b** and **3d** decreased over continued incubation (see 48 h values in Figure 4).




Figure 3: Protein binding of bis-NHC gold(I) complexes **3a** – **3d** to fetal calf serum proteins (left) and bovine serum albumin (right), incubation concentration 3.0 μ M.




Figure 4: Cellular uptake in MCF-7 breast carcinoma cells of bis-NHC complexes **3a** – **3d** with fetal calf serum (top) or without (bottom), incubation concentration 3.0 μM .

Inhibition of TrxRs

The inhibition of mammalian and bacterial TrxRs was evaluated as a possible mode of drug action using purified rat liver TrxR and TrxR from *E. coli* (Table 3). Against rat TrxR, complexes **3a-3d** displayed low efficacy, however, the same structure-activity-relationships as for cytotoxicity were noted (positive effects of the phenyl and bromine substituents). The most active gold NHC complex was **3d** with an IC_{50} value of 16.3 μM . Also against bacterial TrxR from *E. coli* only low activities were observed for **3a-3d**. In this case, complex **3c** was the most active agent with an IC_{50} value of 30.0 μM . Notably, **3c** had triggered the strongest antibacterial effects of the gold NHC complexes in this study. Auranofin showed much higher activity against both forms of TrxR.

Compound	rat TrxR IC ₅₀ [μM]	<i>E. Coli</i> TrxR IC ₅₀ [μM]
Auranofin	0.093 ^{+/-0.009}	0.296 ^{+/-0.069}
3a	127.8 ^{+/-6.1}	410.9 ^{+/-42.1}
3b	71.9 ^{+/-5.4}	57.2 ^{+/-6.2}
3c	30.3 ^{+/-4.6}	30.0 ^{+/-7.1}
3d	16.3 ^{+/-3.2}	96.6 ^{+/-13.4}

Table 3: IC₅₀ values of rat TrxR and *E.coli* TrxR inhibition tests. Standard errors are given as superscripts.

Discussion and Conclusions

The biscarbene complexes **3a-d** displayed high cytotoxic activities in cancer cells (IC₅₀ 0.05 – 5.54 μM), which were most pronounced in case of the phenylimidazole-based and bromine-containing NHC ligands (most active complex **3d**: IC₅₀ 0.05 – 0.18 μM). In comparison to the previously studied monocarbene derivatives (**2a-d**) substantially stronger activity was noted with similar structure-activity-relationships (e.g. positive effect of the phenylimidazole structure; compare IC₅₀ [μM] values for **2a** / **3a**: 10.58 - 16.97 / 3.44 - 5.54, **2b** / **3b**: 5.49 - 12.05 / 0.72-1.51, **2c** / **3c**: 4.46 - 8.13 / 0.09 - 0.18 and **2d** / **3d**: 4.73 - 7.20 / 0.05 - 0.18).¹⁷ The effective antiproliferative effects in this study are overall in good agreement with the reported high potencies of various biscarbene NHC complexes.^{10, 15, 18, 19, 28} Cellular uptake studies provided an explanation for the enhanced activities of the phenylimidazole derived compounds, as complexes **3c** and **3d** caused the highest cellular gold levels. Moreover, in the presence of serum the uptake of various biscarbene compounds appears in general higher than that of the respective monocarbene analogues.^{10, 15, 17} The higher cellular uptake and higher cytotoxic activity are likely the consequence of a larger lipophilicity of **3c** and **3d** compared to **3a** and **3b**.

On the other hand, there was a negative effect of the bromine substituent on the cellular uptake, contrasting the results of the cytotoxicity study. This negative effect of halide substituents on the cellular uptake had also been observed with the analogous monocarbene complexes recently.¹⁷ It can be assumed that the biscarbene complexes of this work are stable in organic solvents, like DMF or DMSO, and water-based matrices, like cell culture medium or buffer. Ongoing stability studies with a related biscarbene gold(I) complex showed that the complex was stable for several days in different solvents (unpublished data). Taken together, this indicates that the intact complexes are taken up into the cells, where they trigger their biological activity.

Complexes **3a-3d** displayed comparably low protein binding (4 – 58 %), and the cellular accumulation efficiency was largely independent from the presence of serum in the cell culture media. With this property biscarbene complexes differ from analogous monocarbene

derivatives, which show high protein binding (80 – 100 %) and in consequence lower cellular concentrations.^{10, 15, 17} Our recent studies confirmed the relevance of an intact gold-NHC fragment for cellular accumulation.¹⁷ Taken together with the low protein binding of biscarbene complexes and their lipophilic / cationic nature, this provides a possible explanation for the generally enhanced cellular bioavailability of gold(I) biscarbene complexes.

Complexes **3a-3d** also inhibited the growth of Gram-positive bacteria, although with general lower potency than the monocarbene analogues **2a-2d** (compare MIC [μ M] values for **2a / 3a**: 2.55 - 2.97 / 79.0 - 100.0, **2b / 3b**: 0.64 - 3.12 / 12.2 - 70.0, **2c / 3c**: 4.16 - 6.45 / 1.7 - 8.7 and **2d / 3d**: 6.25 - 12.51 / 9.6 - 25.5).¹⁷ The preference for Gram-positive over Gram-negative bacteria has also been reported for Auranofin and is likely related to the strong dependence of these bacteria on an intact Trx/TrxR system.²⁹ Experiments on the inhibition of mammalian and bacterial TrxRs afforded moderate to very low activities (IC₅₀ 16.3 – 410.9 μ M), however, the most active complexes **3c** and **3d** were also the most cytotoxic and most efficient antibacterial agents. It should be noted here that much higher activity against TrxRs could be achieved with various monocarbene species (e.g. **2a-d**: IC₅₀ 0.042 – 0.446 μ M), whereas biscarbene gold(I) NHC complexes are usually weaker TrxR inhibitors.^{15, 17, 18} The relevance of TrxR inhibition for the biological activity of **3a-3d** remains elusive at this stage. On the one hand, the excellent accumulation capability of the complexes led to elevated cellular levels that makes a contribution of TrxR inhibition very likely. For example, taking characteristics of MCF-7 cells into account^{30, 31} the value obtained with **3c** after 24h (9.91 nmol/mg cellular protein uptake from serum containing medium, Figure 4 top) corresponds to a molar concentration of 1119 μ M and with this to an 373-fold accumulation compared to the exposure concentration (3.0 μ M). On the other hand, other mechanisms of action (TrxR related and TrxR unrelated) have been identified for various biscarbene gold NHC derivatives, including the targeting of mitochondria,^{9, 15} binding to DNA G-quadruplexes,^{12, 32} interference with the ASK1-p38-MAPK signaling,³³ or antiangiogenic and antivasular effects.¹⁰

The antimicrobial effects of Auranofin and other gold(I) complexes against various microorganisms have recently attracted increasing attention, and studies on the underlying mode of action are of high interest.^{26, 29, 34-40} Regarding the antibacterial effects, the inhibition of bacterial TrxRs by Auranofin has been demonstrated recently and appears to be of major relevance.^{17, 26, 29, 38}

In conclusion, biscarbene gold(I) complexes represent strongly cytotoxic and moderately antibacterial agents. The strong cellular uptake was found to be a crucial factor for obtaining highly antiproliferative agents. In particular, high activities were noted for derivatives containing a phenylimidazole-based NHC ligand. Several types of gold complexes have

demonstrated both antibacterial and cytotoxic effects. The underlying reasons are not completely understood; however, most likely similar modes of action are important in both species (e.g. inhibition of TrxRs). For future therapeutic applications, it will be important to provide structure-activity-relationship data to separate cytotoxic from antibacterial properties.

Acknowledgements

The financial support by Deutsche Forschungsgemeinschaft (DFG) is gratefully acknowledged (project codes: OT338/12-1 and BR3572/4-1). We thank Dr. Raimo Franke (HZI) for fruitful discussions and his support of this study.

Experimental

General

All reagents were obtained from Sigma-Aldrich (Switzerland) or Fluka Analytical. Bovine serum albumin was from Sigma Aldrich, fetal calf serum (FCS) from Biochrom GmbH Berlin, cell culture media were from Thermo Fisher Scientific Bremen, cell lines and bacteria strains were from CLS Cell Lines Service GmbH Eppelheim (RC-124) or Leibniz Institute DSMZ - German Collection of Microorganisms and Cell Cultures GmbH Braunschweig (bacteria, HT-29, MCF-7, MDA-MB-231, tryptic soy broth was from MP Biomedicals GmbH Eschwege, Bacto yeast extract and Bacto agar were from BD Becton Dickinson Heidelberg, Müller-Hinton broth was from Carl Roth GmbH + Co. KG Karlsruhe, D(+) glucose was from Merck KGaA Darmstadt. Tryptic Soy Yeast contains Tryptic Soy Broth (30 g/L) and Bacto Yeast Extract (3g/L). Müller-Hinton-Broth plus Glucose contains Müller-Hinton-Broth (21g/L) and D(+)Glucose (1%). Bacto Agar was used in a concentration of 15 g/L. The purities of the new synthesized compounds were proved by elemental analysis (Flash EA 1112, Thermo Quest) and differed less than 0.5 % from the predicted values. ¹H NMR spectra and ¹³C NMR spectra were recorded using a Bruker AV II-400 or Bruker DRX-400 AS NMR spectrometer. Mass spectra were recorded on an Agilent Technologies 6120 Quadrupole LC/MS (ESI source, positive mode) or an Advion expression^L CMS (ESI source, negative mode). For the absorption measurements in both enzyme assays a Perkin Elmer 2030 Multilabel Reader VICTORTM X4 was used. The amount of gold was detected by HR-CS AAS contrAA-700 from Analytik Jena. Compounds **1a-1d** and **2a-2d** were prepared as described recently.¹⁷

Synthesis

General procedure for synthesis of NHC-Au-NHC⁺ I⁻ complexes **3a-d**

An equivalent amount of 0.23 – 0.34 mmol of the respective (phenyl)imidazolium gold(I) chloride and the corresponding imidazolium iodide were dissolved in 20 mL of a 1:1 mixture of dichloromethane/methanol. After addition of 1.1 equivalents of potassium carbonate the reaction was stirred over 10d at room temperature with light protection. To remove the excess of potassium carbonate solvents were removed, the residue was resuspended in dichloromethane and filtered. The complexes were isolated by evaporating the filtrate and dried under vacuum at 50 °C over a period of 72 h.

[Bis(1,3-diethyl-imidazol-2-ylidene)]gold(I) iodide **3a**

The synthesis of the imidazolium cation of **3a** with different counter ions has been reported^{9, 20}; see general procedure; starting material: Chlorido(1,3-diethyl-imidazol-2-ylidene)gold(I) (120.0 mg, 0.34mmol) and 1,3-diethyl-imidazolium iodide (84.8 mg, 0.34mmol), yield: 141.7 mg (0.25mmol, 74%), white powder; ¹H-NMR (400 MHz, CDCl₃-d₁) δ ppm 7.25 (s, 4H, Im-H4 + Im-H5), 4.31 (q, ³J_{H,H} = 7.3 Hz, 8H, CH₂), 1.55 (t, ³J_{H,H} = 7.4 Hz, 12H, CH₃); ¹³C-NMR (101 MHz, CDCl₃-d₁) δ ppm 182.4 (2C, Im-C₂_{quat.}), 121.2 (4C, Im-C4H + Im-C5H), 46.6 (4C, CH₂), 17.1 (4C, CH₃); elemental analysis: C₁₄H₂₄AuIN₄ (calc. %/found %) C (29.38/29.57), H (4.23/4.17), N (9.79/ 9.64); MS (ESI): m/z 445.2 [M - I]⁺ and m/z 127.0 [I]⁺.

[Bis(4-bromo-1,3-diethyl-imidazol-2-ylidene)]gold(I) iodide **3b**

see general procedure; starting material: Chlorido(4-bromo-1,3-diethyl-imidazol-2-ylidene)gold(I) (120.0 mg, 0.28mmol) and 4-Bromo-1,3-diethyl-imidazolium iodide (91.2 mg, 0.28mmol), yield: 131.5 mg (0.18mmol, 66%), pale yellow powder; ¹H-NMR (400 MHz, CDCl₃-d₁) δ ppm 7.27 (s, 2H, Im-H5), 4.41 (q, ³J_{H,H} = 7.3 Hz, 4H, CH₂), 4.36 (q, ³J_{H,H} = 7.3 Hz, 4H, CH₂), 1.56 (t, ³J_{H,H} = 7.4 Hz, 6H, CH₃), 1.51 (t, ³J_{H,H} = 7.3 Hz, 6H, CH₃); ¹³C-NMR (101 MHz, CDCl₃-d₁) δ ppm 184.2 (2C, Im-C₂_{quat.}), 121.1 (2C, Im-C5H), 105.5 (2C, Im-C4Br_{quat.}), 47.6 (2C, CH₂), 45.7 (2C, CH₂), 16.6 (2C, CH₃), 16.4 (2C, CH₃); elemental analysis: C₁₄H₂₂AuBr₂IN₄ (calc. %/found %) C (23.03/23.50), H (3.04/2.97), N (7.67/ 7.67); MS (ESI): m/z 603.0 [M - I]⁺ and m/z 127.0 [I]⁺.

[Bis(1,3-diethyl-4-phenyl-imidazol-2-ylidene)]gold(I) iodide **3c**

see general procedure; starting material: Chlorido(1,3-diethyl-4-phenyl-imidazol-2-ylidene)]gold(I) (120.0 mg, 0.28mmol) and 1,3-diethyl-4-phenyl-imidazolium iodide (91.0 mg, 0.28mmol), yield: 173.2 mg (0.24mmol, 86%), white powder; ¹H-NMR (400 MHz, CDCl₃-d₁) δ ppm 7.50 (m, 6H, Ph-H_{para/meta}), 7.41 (m, 4H, Ph-H_{ortho}), 7.17 (s, 2H, Im-H5), 4.43 (q, 4H, ³J_{H,H} = 7.3 Hz, CH₂), 4.27 (q, 4H, ³J_{H,H} = 7.3 Hz, CH₂), 1.62 (t, 6H, ³J_{H,H} = 7.2 Hz, CH₃), 1.43 (t, 6H, ³J_{H,H} = 7.2 Hz, CH₃); ¹³C-NMR (101 MHz, CDCl₃-d₁) δ ppm 183.5 (2C, Im-C₂_{quat.}), 135.0 (2C, Im-C₄_{quat.}), 129.7 (2C, Ph-C4H), 129.5 (4C, Ph-C3H + Ph-C5H), 129.1 (4C, Ph-C2H + Ph-

C6H), 127.4 (2C, Ph-C1_{quat.}), 119.0 (2C, Im-C5H), 47.1 (2C, CH₂), 44.0 (2C, CH₂), 17.6 (2C, CH₃), 17.1 (2C, CH₃); elemental analysis: C₂₆H₃₂AuIN₄ (calc. %/found %) C (43.11/43.47), H (4.45/4.34), N (7.73/7.59); MS (ESI): m/z 597.2 [M- I]⁺ and m/z 127.0 [I]⁻.

[Bis(4-(4-bromophenyl)-1,3-diethyl-imidazol-2-ylidene)]gold(I) iodide **3d**

see general procedure; starting material: Chlorido(4-(4-bromophenyl)-1,3-diethyl-imidazol-2-ylidene)]gold(I) (120.0 mg, 0.23mmol) and (4-(4-bromophenyl)-1,3-diethyl-imidazolium iodide (95.5 mg, 0.23mmol), yield: 163.5 mg (0.18mmol, 79%), white powder; ¹H-NMR (400 MHz, CDCl₃-d₇) δ ppm 7.64 (m, 4H, Ph-H2 + Ph-H6), 7.31 (m, 4H, Ph-H3 + Ph-H5), 7.21 (s, 2H, Im-H5), 4.42 (q, 4H, ³J_{H,H} = 7.3 Hz, CH₂), 4.28 (q, 4H, ³J_{H,H} = 7.3 Hz, CH₂), 1.61 (t, 6H, ³J_{H,H} = 7.2 Hz, CH₃), 1.41 (t, 6H, ³J_{H,H} = 7.2 Hz, CH₃); ¹³C-NMR (101 MHz, CDCl₃-d₇) δ ppm 183.9 (2C, Im-C2_{quat.}), 133.7 (2C, Im-C4_{quat.}), 132.4 (4C, Ph-C3H + Ph-C5H), 131.0 (4C, Ph-C2H + Ph-C6H), 126.3 (2C, Ph-C1_{quat.}), 124.2 (2C, Ph-C4Br_{quat.}), 119.4 (2C, Im-C5H), 47.1 (2C, CH₂), 44.2 (2C, CH₂), 17.5 (2C, CH₃), 17.0 (2C, CH₃); elemental analysis: C₂₆H₃₀AuBr₂IN₄ (calc. %/found %) C (35.40/35.71), H (3.43/3.45), N (6.35/6.19); MS (ESI): m/z 755.0 [M- I]⁺ and m/z 127.0 [I]⁻.

Cell culture

HT-29 colon carcinoma cells, MDA-MB-231 breast cancer cells, MCF-7 breast carcinoma cells were maintained in Dulbecco's Modified Eagle Medium (4.5 g/L D-Glucose, L-Glutamine, Pyruvate), which was supplemented with gentamycin (50 mg/L) and fetal bovine serum superior, standardized (Biochrom GmbH, Berlin) (10 % v/v), and were passaged once a week. RC-124 healthy human kidney cells were maintained in McCoy's 5A (modified, with L-Glutamine) medium which was supplemented with gentamycin (50 mg/L) and fetal bovine serum superior, standardized (Biochrom GmbH, Berlin) (10 % v/v), and were also passaged once a week. For experiments with RC-124 cells, microtiter plates had been pretreated in the following way: 30 μL of a sterilized gelatine solution (1.5 % (m/V)) were added to each well of flat bottom 96-well plates, the plates were covered with their lids, incubated for 1h at 37°C, the excess solution was removed, the wells were washed with PBS 7.4 pH, and the new cell-culture medium was added.

Antiproliferative Assay in tumorigenic and non-tumorigenic cells

The antiproliferative effects were determined according to a recently used method with minor modifications. In short: a volume of 100 μL of HT-29 cells (2565 cells/mL), MDA-MB-231 cells (4120 cells/mL), MCF-7 cells (4840 cells/mL) or RC-124 cells (1460 cells/mL) was transferred into the wells of 96-well plates (note: for RC-124 pretreated plates were used, see above) and incubated at 37 °C / 5 % CO₂ for 72 h (MCF-7, MDA-MB-231, RC-124) or

48 h (HT-29). Stock solutions of the compounds in dimethylformamide (DMF) were freshly prepared and diluted with the respective cell culture medium to graded concentrations (final concentration of DMF: 0.1 % v/v). After 72 h (HT-29) or 96 h (MCF-7, MDA-MB-231, RC-124) of exposure, the cell biomass was determined by crystal violet staining and the IC₅₀ value was determined as the concentration that caused 50 % inhibition of cell proliferation compared to an untreated control. Results were calculated as the mean values of three independent experiments.

Apoptosis induction in drug resistant Nalm-6 cells

Measurement of DNA fragmentation in Nalm-6 cells: Apoptotic cell death was determined by a modified cell cycle analysis, which detects DNA fragmentation at the single-cell level. For measurement of DNA fragmentation cells were seeded at a density 1×10^5 cells/mL and treated with different concentrations of **3d**. After 72 h of incubation, cells were collected by centrifugation at 300 g for 5 min, washed with PBS at 4 °C and fixed in PBS/formaldehyde (2 %, v/v) on ice for 30 min. After fixation, cells were incubated with ethanol/PBS (2:1, v/v) for 15 min, pelleted, and resuspended in PBS containing RNase A (40 mg/mL). After incubation for 30 min at 37 °C, cells were pelleted again and finally resuspended in PBS containing propidium iodide (50 mg/mL). Nuclear DNA fragmentation was then quantified by flow cytometric determination of hypodiploid DNA. Data were collected and analyzed by using a FACScan (Becton Dickinson, Heidelberg, Germany) equipped with the CELLQuest software. Data are given in % hypoploidy (subG1), which reflects the number of apoptotic cells.

Antibacterial Screening

Overnight cultures of the bacteria were grown aerobically at 37 °C in Müller Hinton broth with added 1% glucose and pH 7.2 for Gram-negative strains, or with Trypticase soy yeast extract medium (TSY – 30 g/L trypticase soy broth, 3 g/L yeast extract, pH 7.2) for Gram-positive strains. The cultures were adjusted to an OD_{600nm} of 0.001, which resulted in a final start OD_{600nm} of 0.0005 in the test. 25 µL of test culture was added to 25 µL of a serial dilution of the test compounds in the appropriate medium for the different strains in accordance with standardized procedures in 384 well plates. Test compounds from stock solutions in DMF (complexes **3a** – **3d**, Auranofin) DMSO (Linezolid) or water (Ciprofloxacin, Amikacin) were used at final concentrations of 100, 50, 25, 12.5, 6.25, 3.125, 1.56, 0.78, 0.39, 0.2 µM. As positive control compounds, Linezolid (both MRSA strains) Ciprofloxacin (*E. faecium*, *E. coli*, *A. baumannii*, *K. pneumoniae*), Amikacin (*P. aeruginosa*) were applied. The highest DMF/DMSO concentration in the assay was 1%, which had no apparent effect on the growth of the bacteria. After an incubation time of 18 h at 37 °C under moist conditions, the optical density at 600 nm was measured with a Fusion Universal Microplate Analyser (Perkin–

Elmer, Waltham, USA). The lowest concentration that completely suppressed growth defined the MIC values. The MIC values were determined by curve fitting with Sigma Plot. All values represent averages from at least three independent experiments. The following bacterial strains were used. Gram-negative: *Acinetobacter baumannii* (DSM 30007), *Escherichia coli* (DSM 1116), *Klebsiella pneumoniae* (DSM 11678) and *Pseudomonas aeruginosa* PA7 (DSM 24068). Gram-positive: *Enterococcus faecium* (DSM 20477), *Staphylococcus aureus* MRSA (clinical isolate, RKI 11-02670) and *Staphylococcus aureus* MRSA (DSM 11822).

Protein binding studies

The precipitation assay, based on the method of Ma et al., was modified and performed with solutions containing bovine serum albumin or with fetal calf serum.^{17, 41} An aliquot (11 mL) of DMEM cell culture medium was supplemented with FCS (1.1 mL; standardized) or with BSA (440 mg). An aliquot (1.0 mL) of the corresponding solution was used for matrix-matched calibration and was treated like the other samples without incubation. Stock solutions of the test compounds were prepared in DMF (concentration: 3.0 mM). An aliquot (10 mL) of each solution was pipetted into cell culture medium (10 mL) containing either FCS or BSA and carefully mixed (final incubation concentration: 3 μ M). The reaction mixture was incubated at 37 °C for 48 h under shaking. After the requisite duration (0, 1, 4, 8, 24, 48 h), aliquots (250 mL) of each sample were treated with ice-cooled ethanol (500 mL) and stored at -25 °C for 2 h. Thereafter, the samples were centrifuged (964 g at 4 °C for 15 min). An aliquot (350 mL) of the supernatant was separated and stored at -25 °C. The experiment was carried out in duplicate. Aliquots (100 mL) of each sample were treated with 13 % nitric acid (10 mL) for stabilization, and the gold contents were quantified by HR-CS AAS (see below). The bonded moiety was calculated as percentage.

AAS Measurements

For the gold and fluorine measurements a contraAA 700 high-resolution continuum-source atomic absorption spectrometer (Analytik Jena AG) was used. Pure samples of the respective complexes were used as standards and calibration was done in a matrix-matched manner (meaning all samples and standards were adjusted to the same protein concentration by dilution with distilled water).

Triton-X 100 (1 %, 10 μ L) as well as ascorbic acid (1 %, 10 μ L), were added to each standard sample (100 μ L). Samples were injected (25 μ L) into coated standard graphite tubes (Analytik Jena AG) and thermally processed as previously described in more detail.¹⁷ Gold was quantified at a wavelength of 242.79 nm. The mean integrated absorbances of triple injections were used throughout the studies. The final results of gold concentrations were calculated from data obtained in two independent experiments and are expressed as nmol of

metal per mg of cellular protein and as cellular molar concentration [mM]. The procedure for calculating the cellular molar concentration in MCF-7 cell lines is described in the literature.

Cellular Uptake Studies in MCF-7 Cells

The cellular metal uptake was determined according to previously described methods.^{17, 42} In short: MCF-7 breast carcinoma cells were grown until at least 75 - 80 % confluency in 150 cm² cell-culture flasks. Stock solutions of the compounds in DMF were prepared and diluted with cell-culture medium to a final concentration of 3.0 μM immediately before use (final DMF concentration: 0.1 % v/v). The cell culture medium of the flasks was replaced with the medium that contained the metal compound (20 mL) and the flasks were incubated at 37 °C / 5 % CO₂ up to 48 h. After the desired incubation period the uptake was stopped by removing cell culture medium. The cells were washed with PBS (10 mL), the washing solution was removed, and the cells were isolated after 6 min trypsinization (2.4 mL trypsin solution 0.05 %, containing EDTA 0.004 %) by centrifugation (5 min, 1096 g). The obtained cell pellets were stored at -20 °C for further use. For metal and protein quantification the pellets were resuspended in demineralized water (1.0 mL) and lysed 30 min by ultra-sonication. The protein content of lysates was determined by the Bradford method and the metal content was determined by AAS as described below.

Inhibition of mammalian TrxR

To determine the inhibition of mammalian TrxR an established microplate reader based assay was performed.^{14, 17} Commercially available rat liver TrxR (from Sigma-Aldrich) was used and diluted with distilled water to achieve a concentration of 3.58 U/mL. The compounds were freshly dissolved as stock solutions in DMF. 25 μL aliquots of the enzyme solution and 25 μL of either potassium phosphate buffer pH 7.0 containing the compounds in graded concentrations or 25 μL buffer without compounds but DMF (positive control) were added. 50 μL of a blank solution (DMF in buffer) was also prepared (final concentrations of DMF: 0.5 % v/v). The resulting solutions were incubated with moderate shaking for 75 min at 37°C in a 96-well plate. To each well, 225 μL reaction mixture (1 mL reaction mixture consists of 500 μL potassium phosphate buffer pH 7.0, 80 μL EDTA solution (100 mM, pH 7.5), 20 μL BSA solution (0.2 %), 100 μL of NADPH solution (20 mM) and 300 μL distilled water) were added and the reaction started immediately by addition of 25 μL of a 20 mM ethanolic DTNB solution. After proper mixing, the formation of 5-TNB was monitored with a microplate reader at 405 nm 10 times in 35 s intervals for about 6 min. The increase in 5-TNB concentration over time followed a linear trend ($r^2 \geq 0.990$), and the enzymatic activities were calculated as the slopes (increase in absorbance per second) thereof. For each tested compound, the noninterference with the assay components was confirmed by a negative

control experiment using an enzyme-free test solution. The IC₅₀ values were calculated as the concentration of compound decreasing the enzymatic activity of the untreated control by 50 % and are given as the means and error of three repeated experiments.

TrxR *E.coli* Inhibition Assay

The DTNB-coupled thioredoxin reductase inhibition assay for *E. coli* was partly adopted and modified from Lu et al.^{17, 43} Commercially available *E. coli* TrxR and its natural substrate *E. coli* Trx (both from Sigma-Aldrich) were used and diluted with distilled water to achieve a concentration of 35.5 U/mL for the enzyme and 0.77 µg/mL for the substrate. The compounds were freshly dissolved as stock solutions in DMF. 10 µL aliquots of the enzyme solution, 10 µL substrate solution, 100 µL NADPH (200 mM) in TE buffer were mixed in a well with 20 µL TE buffer pH 7.5 (consists of Tris-HCl 50 mM and EDTA 1 mM in aqueous solution) containing the compounds in graded concentrations or 20 µL buffer solution without compounds (control). Blank solutions: 100 µL NADPH (200 mM) and 40 µL of a DMF/buffer mixture were added and the resulting solutions were incubated with moderate shaking for 75 min at 25 °C in a 96-well plate (final concentrations of DMF: 0.5% v/v). To each well, 100 µL of reaction mixture (containing NADPH 200 µM and DTNB 5 mM in TE buffer solution) were added and the reaction started immediately. After proper mixing, the formation of 5-TNB was monitored with a microplate reader (Perkin-Elmer Victor X4) at 405 nm 10 times in 35 s intervals for about 6 min. The values were corrected using the absorbances of the blank solution. The increase in 5-TNB concentration over time followed a linear trend ($r^2 \geq 0.990$), and the enzymatic activities were calculated as the slopes (increase in absorbance per second) thereof. For each tested compound, the noninterference with the assay components was confirmed by a negative control experiment, where the highest test compound concentration is used and the enzyme aliquot solution is replaced by the same amount of TE buffer. The IC₅₀ values were calculated as the concentration of compound decreasing the enzymatic activity of the positive control by 50 % and are given as the means and error of three repeated experiments.

Conflict of Interest

The authors declare no competing interests.

References

1. I. Ott, *Coord. Chem. Rev.*, 2009, **253**, 1670-1681.
2. T. Zou, C. T. Lum, C. N. Lok, J. J. Zhang and C. M. Che, *Chem Soc Rev*, 2015, **44**, 8786-8801.
3. B. Bertrand and A. Casini, *Dalton Trans.*, 2014, **43**, 4209-4219.
4. B. D. Glisic and M. I. Djuran, *Dalton Trans.*, 2014, **43**, 5950-5969.
5. W. Liu and R. Gust, *Coord. Chem. Rev.*, 2016, **329**, 191-213.
6. R. Rubbiani, B. Wahrig and I. Ott, *J. Biol. Inorg. Chem.*, 2014, **19**, 961-965.
7. R. Koch, *Verhandlungen des X. Internationalen Medizinischen Kongresses, Berlin 1890. 1891 Bd.I. Verlag von August Hirschwald, Berlin*, 1890.
8. R. M. Snyder, C. K. Mirabelli and S. T. Crooke, *Biochem. Pharmacol.*, 1986, **35**, 923-932.
9. J. L. Hickey, R. A. Ruhayel, P. J. Barnard, M. V. Baker, S. J. Berners-Price and A. Filipovska, *J. Am. Chem. Soc.*, 2008, **130**, 12570-12571.
10. J. K. Muenzner, B. Biersack, A. Albrecht, T. Rehm, U. Lacher, W. Milius, A. Casini, J. J. Zhang, I. Ott, V. Brabec, O. Stuchlikova, I. C. Andronache, D. Schuppan, L. Kaps and R. Schobert, *Chemistry*, 2016, **22**, 18953 – 18962.
11. B. Bertrand, A. Citta, I. L. Franken, M. Picquet, A. Folda, V. Scalcon, M. P. Rigobello, P. Le Gendre, A. Casini and E. Bodio, *J Biol Inorg Chem*, 2015, **20**, 1005-1020.
12. B. Bertrand, L. Stefan, M. Pirrotta, D. Monchaud, E. Bodio, P. Richard, P. Le Gendre, E. Warmerdam, M. H. de Jager, G. M. Groothuis, M. Picquet and A. Casini, *Inorg Chem*, 2014, **53**, 2296-2303.
13. R. Rubbiani, L. Salassa, A. de Almeida, A. Casini and I. Ott, *ChemMedChem*, 2014, **9**, 1205-1210.
14. R. Rubbiani, I. Kitanovic, H. Alborzinia, S. Can, A. Kitanovic, L. A. Onambele, M. Stefanopoulou, Y. Geldmacher, W. S. Sheldrick, G. Wolber, A. Prokop, S. Wolf and I. Ott, *J. Med. Chem.*, 2010, **53**, 8608-8618.
15. R. Rubbiani, S. Can, I. Kitanovic, H. Alborzinia, M. Stefanopoulou, M. Kokoschka, S. Mönchgesang, W. S. Sheldrick, S. Wölfl and I. Ott, *J. Med. Chem.*, 2011, **54**, 8646-8657.
16. R. Rubbiani, E. Schuh, A. Meyer, J. Lemke, J. Wimberg, N. Metzler-Nolte, F. Meyer, F. Mohr and I. Ott, *Med. Chem. Commun.*, 2013, **4**, 942-948.
17. C. Schmidt, B. Karge, R. Misgeld, A. Prokop, R. Franke, M. Bronstrup and I. Ott, *Chemistry*, 2017, **23**, 1869-1880.
18. W. Liu, K. Bendorf, M. Proetto, A. Hagenbach, U. Abram and R. Gust, *J. Med. Chem.*, 2012, **55**, 3713-3724.

19. J. K. Muenzner, B. Biersack, H. Kalie, I. C. Andronache, L. Kaps, D. Schuppan, F. Sasse and R. Schobert, *ChemMedChem*, 2014, **9**, 1195-1204.
20. M. Baker, P. J. Barnard, S. J. Berners-Price, S. K. Brayshaw, J. L. Hickey, B. W. Skelton and A. H. White, *Dalton Trans.*, 2006, 3708-3715.
21. T. Zou, C. T. Lum, S. S.-Y. Chui and C.-M. Che, *Angewandte Chemie*, 2013, **52**, 1-5.
22. T. Zou, C. T. Lum, C.-N. Lok, W.-P. To, K.-H. Low and C.-M. Che, *Angew Chem Int Ed*, 2014, **53**, 5810-5814.
23. J. Fernandez-Gallardo, B. T. Elie, M. Sanau and M. Contel, *Chem Commun (Camb)*, 2016, **52**, 3155-3158.
24. T. V. Serebryanskaya, A. A. Zolotarev and I. Ott, *Med. Chem. Commun.*, 2015, **6**, 1186-1189.
25. C. V. Maftai, E. Fodor, P. G. Jones, M. Freytag, M. H. Franz, G. Kelter, H. H. Fiebig, M. Tamm and I. Neda, *Eur J Med Chem*, 2015, **101**, 431-441.
26. J. P. Owings, N. N. McNair, Y. F. Mui, T. N. Gustafsson, A. Holmgren, M. Contel, J. B. Goldberg and J. R. Mead, *FEMS Microbiol Lett*, 2016, **363**.
27. L. Kaps, B. Biersack, H. Muller-Bunz, K. Mahal, J. Munzner, M. Tacke, T. Mueller and R. Schobert, *J Inorg Biochem*, 2012, **106**, 52-58.
28. J. F. Arambula, R. McCall, K. J. Sidoran, D. Magda, N. A. Mitchell, C. W. Bielawski, V. M. Lynch, J. L. Sessler and K. Arumugam, *Chem Sci*, 2016, **7**, 1245-1256.
29. M. B. Harbut, C. Vilcheze, X. Luo, M. E. Hensler, H. Guo, B. Yang, A. K. Chatterjee, V. Nizet, W. R. Jacobs, Jr., P. G. Schultz and F. Wang, *Proc. Natl. Acad. Sci. U S A*, 2015, **112**, 4453-4458.
30. R. Gust, B. Schnurr, R. Krauser, G. Bernhardt, M. Koch, B. Schmid, E. Hummel and H. Schönenberger, *J. Cancer Res. Clin. Oncol.*, 1998, **124**, 585-597.
31. H. Reile, G. Bernhardt, M. Koch, H. Schönenberger, M. Hollstein and F. Lux, *Cancer Chemother Pharmacol*, 1992, **30**, 113-122.
32. C. Bazzicalupi, M. Ferraroni, F. Papi, L. Massai, B. Bertrand, L. Messori, P. Gratteri and A. Casini, *Angew Chem Int Ed Engl*, 2016, **55**, 4256-4259.
33. X. Cheng, P. Holenya, S. Can, H. Alborzinia, R. Rubbiani, I. Ott and S. Wolf, *Mol. Cancer*, 2014, **13**, 221.
34. F. Angelucci, A. A. Sayed, D. L. Williams, G. Boumis, M. Brunori, D. Dimastrogiovanni, A. E. Miele, F. Pauly and A. Bellelli, *J Biol Chem*, 2009, **284**, 28977-28985.
35. S. Jackson-Rosario, D. Cowart, A. Myers, R. Tarrien, R. L. Levine, R. A. Scott and W. T. Self, *J Biol Inorg Chem*, 2009, **14**, 507-519.
36. Y. Hokai, B. Jurkowicz, J. Fernandez-Gallardo, N. Zakirkhodjaev, M. Sanau, T. R. Muth and M. Contel, *J Inorg Biochem*, 2014, **138**, 81-88.

37. T. Gamberi, T. Fiaschi, A. Modesti, L. Massai, L. Messori, M. Balzi and F. Magherini, *Int J Biochem Cell Biol*, 2015, **65**, 61-71.
38. A. Debnath, D. Parsonage, R. M. Andrade, C. He, E. R. Cobo, K. Hirata, S. Chen, G. Garcia-Rivera, E. Orozco, M. B. Martinez, S. S. Gunatilleke, A. M. Barrios, M. R. Arkin, L. B. Poole, J. H. McKerrow and S. L. Reed, *Nat. Med.*, 2012, **18**, 956-960.
39. E. R. Sharlow, S. Leimgruber, S. Murray, A. Lira, R. J. Sciotti, M. Hickman, T. Hudson, S. Leed, D. Caridha, A. M. Barrios, D. Close, M. Grogl and J. S. Lazo, *ACS Chem. Biol.*, 2014, **9**, 663-672.
40. M. I. Cassetta, T. Marzo, S. Fallani, A. Novelli and L. Messori, *Biomaterials*, 2014, **27**, 787-791.
41. J. Ma, G. Stoter, J. Verweij and J. H. Schellens, *Cancer Chemother Pharmacol*, 1996, **38**, 391-394.
42. H. Scheffler, Y. You and I. Ott, *Polyhedron*, 2010, **29**, 66-69.
43. J. Lu, A. Vlamis-Gardikas, K. Kandasamy, R. Zhao, T. N. Gustafsson, L. Engstrand, S. Hoffner, L. Engman and A. Holmgren, *FASEB J*, 2013, **27**, 1394-1403.

Antiproliferative effects against human cells

HT-29 / MCF-7 / MDA-MB-231
RC-124

